

## EYELID SURGERY

### Pre-Operative and Post-Operative Instructions

#### PREPARING FOR SURGERY

##### One month before surgery:

- See your primary care provider within 30 days of your surgery for your pre-operative exam and ensure that they send Dr. Ridgway's office a copy of your medical clearance note. F:406-587-2120, E:office@ridgwayplasticsurgery.com. Medical clearance will not be requested for every patient.
- Stop tobacco use and avoid being around anyone who smokes for at least 1 month before and 2 weeks after surgery. This includes nicotine patches and gum, chewing tobacco and vaping. Smoking impairs wound healing and increases your risk of infection. It also increases your risk of anesthesia complications including pneumonia, cardiac events and blood clots. Due to a strong risk for delayed healing, we will perform a urine test prior to your surgery to test for by-products of smoking. If the test is positive, your surgery will be cancelled. If you have not been able to stop, please let us know as soon as possible and we can reschedule your surgery.
- Start eating a healthy diet to prepare your body to heal smoothly. Increase your protein intake to ideally 70 grams per day and add foods high in Vitamin C to your diet -- dark leafy greens with kale and spinach, strawberries, broccoli, bell peppers, kiwi, cantaloupe, Brussels sprouts, papaya and oranges. You may also want to supplement with a daily zinc supplement or multivitamin. Zinc is found in shrimp, cashews, lentils, sesame and pumpkin seeds, lamb, turkey and quinoa. Protein, Vitamin C and zinc have all been shown to help healing.
- If you have diabetes, you are at increased risk for wound infection and problems healing. We want to have your glucose levels optimized and well controlled prior to surgery. Work with your PCP to ensure this goal is reached.
- If you have rheumatoid arthritis, ulcerative colitis, lupus or are taking any chemotherapy, steroids or immunosuppression for another reason, you are at increased risk for wound healing problems and infection. Please alert us of these medications as we will attempt to work with your prescribing doctor to stop these medications. If it is not possible to stop these medications, the risks of complications may outweigh the benefits of surgery.
- If you have a history of a blood clot - deep vein thrombosis (DVT), pulmonary embolism (PE), multiple miscarriages, a family history of blood clots - you are at increased risk for a DVT or PE happening again. A PE can be fatal. We will need to speak with you, to your doctor and a hematologist about perioperative blood thinners such as Lovenox®.

##### Two weeks before surgery:

- Come to your pre-operative visit.
- Fill your prescriptions
- If you take blood thinners, such as Coumadin®, Plavix® or Lovenox®, please speak with Dr. Ridgway and your primary care provider about stopping these medications. Sometimes the risks of stopping these medications outweigh the benefits of surgery.

- Fill your prescriptions.
- Stop taking all aspirin and ibuprofen products 14 days before surgery and at least 2 days after your surgery, unless otherwise directed. These products thin your blood and may increase bleeding. To be safe, please take acetaminophen (Tylenol®) only for pain relief in the 2 weeks prior to surgery.

Other bleeding thinning products to avoid are:

- Vitamin E
  - Fish oil
  - Ginkgo biloba
  - Garlic supplements
  - St. John's wort
  - Herbal teas
  - Ginger supplements
  - Excessive intake of avocado, cayenne pepper, cranberries, raisins or dried cherries
- Continue to take your prescribed medications unless otherwise advised by your doctor. If you have questions about your medications, ask your doctor or pharmacist.
  - Stop drinking alcohol at least 7 days prior to surgery. Alcohol consumption (even weekend binge drinking) can impair your ability to stop bleeding and to fight an infection. Alcohol consumption also increases your risk of complications from anesthesia.
  - Stay well hydrated.
  - Arrange for a support person to help you after surgery with the following:
    - Receive instructions for your care
    - Drive you home and stay with you for at least the first 24 hours after surgery
    - Bring you back to the clinic for your first post-op visit

#### The day before your surgery:

- A representative from the office will call you before your surgery to let you know the time of your surgery.
- If you are having general anesthesia, do not eat or drink anything after midnight. Hydrate well the day and week before surgery.
- Pick up your prescriptions.
- Shampoo your hair. Cleanse your face and neck the night before and the morning of your surgery with an antibacterial soap such as Dial® or Lever 2000® or Hibiclens.
- DO NOT use hair spray, make-up, creams, or lotions the day of surgery.
- DO NOT wear contact lenses the day of surgery.
- DO NOT wear any jewelry or nail polish. Sculptured nails may have to be clipped for monitoring purposes.
- You may wish to bring a hat, scarf, and/or sunglasses to wear home following your surgery.

#### Helpful preparation hints:

- Make sure items like milk, juice, and water are in small containers. Gallon and half gallon containers will be too heavy for you to lift.
- Have pre-made dinners on hand so you can cook them in the microwave.
- Prepare foods high in protein such as meat, poultry, fish, and eggs. Eating more protein will help you heal at a faster rate, as will adding dark green vegetables.

- Have crackers or pretzels on hand to eat with your medication or if you have an upset stomach.
- Use paper plates and plastic cups so you do not need to wash dishes.
- Bring items down from high shelves so you do not have to reach. Bring up items on low shelves so you do not need to bend down.
- Have your night clothes, slippers, dressing supplies, and medications set up and ready for when you come home. Have a phone in easy reach, as well as remote controls, and reading material.
- Try to have someone available to run to the store for you. Driving can be hazardous the first two weeks and while you are taking narcotics. Your range of motion may be limited and you may not be able to react quickly enough in response to a dog running into the road or a car cutting you off.

Items to have on hand:

Some supplies which may be helpful during your recovery are:

- Saline drops are for during the day; Lacri-Lube® to use at night to moisten your eyes
- Arnica ointment to apply to areas of bruising
- A gentle laxative for constipation if taking narcotics
- Yogurt or probiotics to prevent yeast infections if on antibiotics
- Thermometer
- Slip on shoes
- Several house robes
- Extra pillows to sleep with your head elevated
- High protein foods
- Antibacterial soap
- Aquaphor® ointment
- Extra gauze or nursing pads
- Paper tape
- A flexible straw for drinking
- A list of phone numbers/emergency contacts for your helpers
- Frozen peas or cooling packs to place over your eyes. (Remember to use a washcloth in between the ice and your skin always.)

#### DAY OF SURGERY

- Arrive at the location for your surgery at the time requested.
- Do not drink or eat anything the morning of your surgery if you are having general anesthesia.
- Take your normal medications.
- Your support person can wait in the surgery waiting room. He or she should plan on waiting about 4-6 hours.
- Your surgery normally takes around 1-2 hours.

The first night of surgery, you are at risk for bleeding under the skin. This would present with increased swelling, increased pain and possibly increased bruising. This may be a surgical emergency so please call and we will ask you to meet us in the clinic or hospital to evaluate you.

#### THE HEALING PROCESS:

During the day, please apply saline drops to your eyes if they feel dry.

Before going to sleep, please apply Lacri-Lube® to your eyes to prevent dryness.

Please sleep with your head elevated above your heart for two weeks.

Apply arnica ointment around your eyes to decrease bruising, if you wish. Do not apply to incisions.

During the first 48 hours, keep iced gauze or ice packs on your eyes to decrease swelling.

### Incision care:

Your incisions will be closed with sutures. They will be visible to you. Bacitracin will be placed over the incisions. The healing process after facial surgery varies with each person. You should expect to feel tired for the first 2 to 3 weeks due to the anesthesia and healing process. Rest often during the day and get a good night's sleep.

- If you have stitches they will be removed in approximately 5 days.
- Swelling is normal, as is mild bruising. If one side is swelling more than the other or the swelling is painful, it could be a fluid collection. Please call to be seen in clinic immediately.
- Expect redness at your incision line for several weeks as your stitches absorb. If the redness spreads away from the incision, please call the clinic.
- Crustiness at the incision site is normal during the first 2 weeks. Please use a Q-Tip to clean off any crusty discharge. Reapply Bacitracin twice daily until the sutures are removed. After the sutures are removed, please continue to clean with Q-Tips and apply Aquaphor® twice daily
- You may shower in 2 days. Allow water and soap to run over the incisions and gently pat them dry.
- Your incisions will get red and bumpy in the first 3 months as your body builds scar tissue. Scar massage will help the scars soften and fade. You may begin scar massage at 6 weeks post surgery (see Scar Massage handout). By your 6-month appointment the scars will have lightened and flattened. However, it takes at least a year for your scars to heal and scar appearance will continue to improve for years.

### Sleeping: Try to get a good night's sleep

Stay in bed, on your back, with your head and shoulders elevated above the heart for 24 hours after your surgery. Your surgeon may ask you to sleep on your back for 2 weeks. Keep your neck in a neutral position with your chin up. Here are some suggestions for a good night's sleep.

- Try sleeping in your recliner chair.
- Have extra pillows in your bed for support: two along your sides and one under your knees to keep you on your back and to relieve lower back pressure. Buy a large "body pillow" or a pillow with arm rests for sitting up in bed.

### Pain Medications:

With any surgery there is some discomfort or pain. We recommend Tylenol® use only for the first two days. After two days, you may also take ibuprofen.

### Showering and bathing

- You may shower 2 days following surgery. Wash by allowing the water and soap to run over your incisions. Pat the incisions dry. You do not want your incisions to be covered with moist dressings.
- Use a Q-Tip to remove dried blood or crusting. Reapply Aquaphor® after showering.

- Be sure to have someone nearby when taking your first shower in case you feel dizzy. Remember your eyes may be swollen.
- Do not soak, swim or hot tub until all incisions are completely healed.

#### Do's and Don't's for the next 6 weeks:

- Keep your head above your heart. Do not bend to lift things.
- DO NOT drive a motor vehicle for 1 week until you can see well and can handle the steering wheel without discomfort. Do not drive while taking any narcotics.
- DO NOT engage in sexual activity for at least 1 week.
- DO NOT smoke or be around anyone who smokes for 2 weeks. Smoking delays healing and can lead to infection.
- DO NOT lift more than 5 pounds or bend at the waist to lift for 2 weeks.
- DO NOT participate in strenuous activities such as running or aerobics for 6 weeks. When you do resume these activities, do so slowly. If it hurts, it is too early.
- Stay out of the sun for the first week and DO NOT tan at the incision lines for at least 12 months. Apply sunscreen to incisions if you are in the sun.
- DO NOT wear makeup until the sutures are removed and incisions healed.
- Do resume walking at a gentle pace.
- You may return to work in 1 to 6 weeks (average time is 2 weeks) depending upon your work activity.
- Do report signs of infection. We want to see you as soon as possible if you have any of these signs.

#### Signs of Infection:

- A temperature over 101°F or 38°C.
- Redness at the incision line that is beginning to spread away from the incision after the first 48 hours.
- Yellow pus-like or foul smelling drainage larger than dime size from the incision or drain sites.
- Increased pain/discomfort that is not relieved by your pain medicine.
- Increased swelling.

#### All surgery has some risk:

Eyelid surgery is not a simple surgery, but is normally safe when performed by an experienced and board-certified plastic surgeon.

There are, however, always risks of complications, including unfavorable scars, seroma or hematoma (fluid collection), difficulty closing your eyes (lagophthalmos), eyelid drooping, dry eye, wound healing problems including skin loss, infection, temporary or permanent loss of sensation or paralysis, asymmetry, chronic pain or complications from anesthesia. These problems are more common in overweight, diabetic and smoking patients.

Eyelid surgery does leave permanent scars. However, the scars will continue to improve for years.

#### CONTACT INFORMATION

Office hours: 8:00 am – 4:00 pm, Monday – Thursday; 8:00 am – 1:00 pm Friday

Please call 406-586-2620 with any questions or concerns.